

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29066 A2

- (51) International Patent Classification⁷: C07K 5/00 (74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi S.p.A., Corso dei Porta Vittoria, 9, I-20122 Milano (IT).
- (21) International Application Number: PCT/EP00/10181
- (22) International Filing Date: 17 October 2000 (17.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
FI99A000217 21 October 1999 (21.10.1999) IT
- (71) Applicant (for all designated States except US): MENARINI RICERCHE S.P.A. [IT/IT]; Via Tito Speri, 10, I-00040 Pomezia (IT).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): PERROTTA, Enzo [IT/IT]; Via Ragazzi del '99, 76, I-50141 Firenze (IT). GIANNOTTI, Danilo [IT/IT]; Via Roma, 128, I-55011 Altopascio (IT). ALTAMURA, Maria [IT/IT]; Viale Matteotti, 54, I-50132 Firenze (IT). GIULIANI, Sandro [IT/IT]; Via Carlo Borroni, 11, I-50012 Bagno a Ripoli (IT). MAGGI, Carlo, Alberto [IT/IT]; Via Michelazzi, 43, I-50141 Firenze (IT).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BASIC MONOCYCLIC COMPOUNDS HAVING NK2 ANTAGONIST ACTION, PROCESSES FOR THEIR PREPARATION, AND FORMULATIONS CONTAINING THEM



(I)

(57) Abstract: Basic monocyclic compounds having general formula (I) useful as NK-2 antagonists are described, processes for their preparation and pharmaceutical compositions containing them are also described.

BEST AVAILABLE COPY

BEST AVAILABLE COPY


WO 01/29066 A2

THIS PAGE BLANK (USPTO)

BASIC MONOCYCLIC COMPOUNDS HAVING NK2 ANTAGONIST ACTION,
PROCESSES FOR THEIR PREPARATION, AND FORMULATIONS
CONTAINING THEM

Scope of invention

5 The present invention refers to new compounds having general formula:



in which:

X may be CH₂, O, S, or SO;

Y is $\text{CH}_2\text{-CO-NH}$ or CO :

20 R and R₁, same or different from one another, represent an H or halogen group, including the respective diastereoisomers and their mixtures, with the proviso that: when Y is CH₂-CO-NH, X is CH₂, and R and R₁ are H, then the stereochemistry of the carbon atom linked to Y is of the R type.

The presence of free amine groups bestows on the compounds specifically basic characteristics, but included in the present invention are also pharmaceutically acceptable salts of the compounds of general formula (I) with organic and inorganic acids chosen in the group consisting of: hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, carbonic acid, acetic acid, trifluoroacetic acid, trichloroacetic acid oxalic acid, malonic acid, malic acid, succinic acid, tartaric acid, citric acid, methanesulphonic acid, and p-toluenesulphonic acid.

30 State of the art

The NK2 receptor of tachykinins is widely expressed in the peripheral nervous system of mammals. One of the various effects produced by selective stimulation


of the NK2 receptor is the contraction of smooth muscle. Hence, NK2-receptor antagonists may be considered agents that are capable of controlling excessive contraction of the smooth muscle in any pathological condition in which release of tachykinins concurs in the genesis of the corresponding disorder. In particular, the bronchospastic component of asthma, coughing, pulmonary irritations, intestinal spasms or local spasms of the bladder and the ureter during cystitis, renal infections and colics may be considered conditions in which the administration of NK2 antagonists may be effective (E.M. Kudiacz *et al.*, Eur. J. Pharmacol., 1993, 36, 17-25).

- 10 Cyclic compounds, in particular cyclic hexapeptides (A.T. McKnight *et al.*, Br. J. Pharmacol., 1991, 104, 355) and bicyclic hexapeptides (V. Pavone *et al.*, WO 93/212227), or cyclic pseudopeptides (L. Quartara *et al.*, J. Med. Chem., 1994, 37, 3630; S.L. Harbeson *et al.*, Peptides, Chemistry and Biology. Proceedings of the Twelfth American Peptide Symposium, 1992, 124) are known in the literature for their high antagonistic activity towards the NK2 receptor of tachykinins.

15 Recently it has been demonstrated in WO 9834949 that monocyclic products having a lower molecular weight and containing only four bifunctional residues bound together with a peptide or pseudopeptide bond present a pharmacological activity equal to or higher than that of known compounds; they are moreover endowed with considerable selectivity in regard to the human NK2 receptor and are consequently proposed as valid alternatives.

20 In the compounds according to the present invention, the presence in the products claimed of a grouping of Formula (II)

25



(II)

bestows on the compounds of Formula (I) a surprising activity towards the NK2 receptor when compared to the closest compounds, namely those described in WO 9834949.

Detailed description of the invention

The purpose of the present invention is therefore to make available new monocyclic compounds containing four residues bound together with a peptide bond and having an antagonistic action on the NK2 receptor, of general formula (I) as previously defined.

- 5 Forming part of the present invention are also the pharmaceutically acceptable salts, the processes for their preparation and the pharmaceutical compositions containing them.

More specifically, the halogen group defined for the substituents R and R₁ may be chosen from among fluorine, chlorine, bromine, and iodine.

- 10 Preferred compounds of the invention are the compounds of general formula (I) in which:

R and R₁, same or different from one another, are chosen in the group consisting of H or F, and the other substituents are as defined above.

The following constitute a group of particularly preferred compounds:

- 15 1) cyclo{-Suc[1-(R)-2(1,4')-bipiperidiny-1'-yl-acetyl amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = CH₂)
- 2) cyclo{-Suc[1-(R)-2(4-morpholin-4-yl-piperidin-1-yl)-acetyl amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = O)
- 3) cyclo{-Suc[1-(R)-2(1,4')-bipiperidiny-1'-yl-acetyl amino]-Trp-Phe-(4-F)-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = F; R₁ = H; Y = CH₂CONH; X = CH₂)
- 20 4) cyclo{-Suc[1-(R)-2(4-morpholinyl-4-piperidiny-1-yl)-acetyl amino]-Trp-(5-F)-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = F; Y = CH₂CONH; X = O)
- 5) cyclo{-Suc[1-(R)-2(4-(1-oxo-1-thiomorpholin-4-piperidin-1-yl)-acetyl amino)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = SO)
- 25 6) cyclo{-Suc[1-(R)-2(4-(1-thiomorpholin-4-yl-piperidin-1-yl)-acetyl amino)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = S)
- 7) cyclo{-Suc[1-(1,4')-bipiperidiny-1'-carbonyl]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (slow-moving) (R = H; R₁ = H; Y = CO; X = CH₂)
- 8) cyclo{-Suc[1-(1,4')-bipiperidiny-1'-carbonyl]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (fast-moving) (R = H; R₁ = H; Y = CO; X = CH₂).
- 30

Pharmaceutically acceptable salts of compounds of Formula (I) include salts with inorganic acids (such as hydrochloric acid, hydrobromic acid, carbonic acid,

sulphuric acid, and phosphoric acid) and organic acids (such as acetic acid, trifluoroacetic acid, trichloro-acetic acid, oxalic acid, malonic acid, malic acid, succinic acid, tartaric acid, citric acid, methanesulphonic acid, and p-toluenesulphonic acid).


- 5 According to the invention, the compounds of Formula (I) containing peptide or pseudopeptide bonds may be obtained by condensation using techniques that are known in the literature.

As may be seen from Formula (I), for each compound two diastereoisomers are possible (the carbon atom to which the Y group is bound being quaternary).

- 10 Consequently, the invention refers to the different diastereoisomers of Formula (I) and their mixtures, apart from the case expressly excluded.


- The products according to the present invention may be prepared applying the synthesis described in WO 9834949 (see also the literature quoted therein) according to which the monocyclic is obtained by condensing, according to
15 techniques that are known in the chemistry of peptides, firstly the two amino-acid residues, and then by adding the diamine system and the dicarboxylic system, and finally by cyclization.

- As a possible alternative to the above synthesis, the products of the present invention may be prepared starting from the diamine system of (R)-1-benzyl-2-(N-benzyloxycarbonylamino)-ethyl amine, to which, according to methodologies
20 known in the chemistry of peptides, Phe, Trp (possibly substituted, respectively by an R group and an R₁ group, in which said groups have the meanings previously defined) and a dicarboxylic system (a derivative of succinic acid) are combined. The linear product obtained is cyclized to obtain the desired monocyclic system
25 having general formula (III)



10 in which Y_1 may be $-NH_2$ or $-COOH$, and R and R_1 have the meanings described above.

The compounds of Formula (III) are then combined with the compounds of Formula (IV)



in which Y_2 may be H or CH_2-COOH , and X has the meaning previously defined, to obtain the final compounds of Formula (I).

20 The compounds of Formula (I) as specified above have been found to be powerful antagonists of the NK_2 receptor of tachykinins, and consequently can be administered as agents capable of controlling the excessive contraction of smooth muscle in any pathological condition in which release of tachykinins concurs in the genesis of the corresponding disorder.

25 In particular, the bronchospastic component of asthma, coughing, pulmonary irritations, intestinal spasms or local spasms of the bladder and the ureter during cystitis, renal infections and colics may be considered conditions in which the administration of the compounds of Formula (I), as NK_2 antagonists, may be effective.

30 The compounds of Formula (I) that form the subject of the present invention or their pharmaceutically acceptable salts are suitable for administration for therapeutic purposes to the higher animals and to man through the parenteral,

oral, inhalational, or sublingual routes, achieving pharmacological effects in accordance with the properties described above. For parenteral (intravenous, intramuscular, and intracutaneous) administration, sterile solutions or lyophilized preparations are used. For nasal instillation and for inhalational and sublingual administration, aqueous solutions, aerosol preparations, powders or capsules are used as required.

The doses of active principle in the above-mentioned compositions may be such to allow the delivery of between 0.1 and 10 mg per kg of body weight.

The following are non-limiting examples of the present invention:

EXAMPLE 1: cyclo{-Suc[1-(R)-2(1,4')-bipiperidiny-1'-yl-acetyl amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(a compound of general formula (I) in which R = H, R₁ = H, Y = CH₂CONH, X = CH₂, and in which C-Y has R configuration)

Used as a starting product is the compound cyclo{-Suc[1-(R)-amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}, referred to as Compound A (compound of Formula (III) in which R = H, R₁ = H, Y₁ = NH₂, and in which the C-Y₁ carbon has R configuration), prepared as follows:

a) Synthesis of (R)-1-benzyl-2(N-benzyloxycarbonylamino) ethyl amine

(R)-1-benzyl-1-(N-*tert*-butyloxycarbonylamino)ethyl amine, obtained as described in G. Kokotos *et al.*, J. Chem. Research (S), 1992, 391, was converted into the corresponding (R)-1-benzyl-1-(N-*tert*-butyloxycarbonylamino)-2-(benzyloxycarbonylamino)ethyl amine, and the latter into (R)-1-benzyl-2-(N-benzyloxycarbonylamino)ethyl amine according to the customary methods of protection and de-protection of amino acids.

b) Synthesis of Boc-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z]

To a solution of Boc-Phe-OH (5.1 g) in 200 ml of anhydrous THF at 5°C were added in order HOBt (4.2 g), EDC.HCl (4.5 g), and finally, drop by drop, a solution of (R)-1-benzyl-2-(N-benzyloxycarbonylamino)ethyl amine (5.5 g) in 30 ml of anhydrous THF. The solution was left overnight under stirring at room temperature. The solvent was evaporated, and the residue was treated with a 5% aqueous solution of KHSO₄. The resulting solid was filtered, washed with water, then with a 10% aqueous solution of NaHCO₃, and finally with a saturated solution

of NaCl up to pH 7. After vacuum drying, 7.2 grams of a colourless solid product were obtained.

MS (ES⁺) : [MH⁺] = 532; HPLC (Method A1): rt = 18.8 min

c) Synthesis of H-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z]

- 5 TFA (25 ml) was added, under stirring at 0°C, to a suspension of the above compound (4.0 g) in CH₂Cl₂ (25 ml). The reaction mixture was kept under stirring for 2 hours at room temperature, and the disappearance of the precursor was monitored by means of HPLC analysis. The solution was dried, the residue was dissolved in AcOEt (100 ml), and the resulting organic phase was washed with a
10 saturated aqueous solution of K₂CO₃ (25 ml), then with brine up to pH 7 (4 x 50 ml), and finally was dried on anhydrous sodium sulphate. By evaporation of the solvent, 3.5 g of a white solid were obtained.

d) Synthesis of Boc-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z]

- HOBT (3.0 g) and EDC.HCl (1.5 g) were added to a solution of Boc-Trp-OH (2.2 g)
15 in THF (100 ml). After stirring for 15 minutes, a solution of H-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z] (3.1 g) in THF (15 ml) was added. The mixture was left under stirring at room temperature overnight; then the solvent was evaporated, and the residue was treated with a 5% aqueous solution of KHSO₄ (100 ml) to obtain a white solid. The solid was filtered and washed again with a 5% aqueous solution of
20 KHSO₄ (2 x 100 ml), then with a 5% aqueous solution of NaHCO₃ (3 x 50 ml), and finally with water (3 x 50 ml), and was then vacuum dried to obtain 4.9 grams of a white solid.

HPLC (Method A3): rt = 18.0 min; MS (TS⁺) : [MH⁺] = 718

e) Synthesis of H-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z]

- 25 TFA (15 ml) was added, under stirring at 0°C, to a suspension of the previous compound (1.0 g) in CH₂Cl₂ (25 ml). The reaction mixture was kept under stirring for 30 minutes at 0°C and for 2 hours at room temperature, and the disappearance of the precursor was monitored by means of HPLC analysis. After evaporation of the solvent, the residue was diluted with AcOEt (100 ml), washed with a 5%
30 aqueous solution of NaHCO₃ (2 x 30 ml) and brine (30 ml).

The organic phase was dried with MgSO₄ and vacuum evaporated at 30°C to obtain 650 mg of the desired compound.

f) Synthesis of Boc-(D)-Asp{Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH-Z]}-OBzl

To a solution of Boc-(D)-Asp-OBzl (690 mg), HOBt (850 mg), and EDCI.HCl (450 mg) in anhydrous DMF (50 ml) was added, under stirring at room temperature and under nitrogen, a solution of the compound of Example 1(e) (1.3 g). The reaction mixture was left under stirring at room temperature for four hours. After evaporation of the solvent in vacuum conditions, the residue was treated with a 5% aqueous solution of KHSO₄ to obtain a solid which was filtered, washed with a 5% aqueous solution of NaHCO₃, with water, and dried. The product was crystallized using ethanol to obtain 850 mg of the desired compound as a white solid.

MS (ES⁺) : [MH⁺] = 923; HPLC (Method A1): rt = 21.1 min

g) Synthesis of Boc-(D)-Asp{Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH₂]}-OH

The compound of Example 1(f) (800 mg) was solubilized in DMF (10 ml) and diluted with MeOH (40 ml), then hydrogenated in the presence of 10% Pd/C (100 mg) at atmospheric pressure and room temperature for 5 hours. The catalyst was filtered and washed with MeOH. After evaporation of the solvent, 500 mg were obtained of the desired compound as a white solid.

MS (ES⁺) : [MH⁺] = 699; HPLC (Method A2): rt = 10.4 min

h) Synthesis of cyclo{-Suc[1(R)NHBoc]-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]}-

To a solution of the compound of Example 1(h) (800 mg) in anhydrous DMF (200 ml) were added, under stirring and in a nitrogen atmosphere, 465 mg of HOBt and 224 mg of EDCI.HCl. The reaction mixture was left under stirring for 5 hours and then, after evaporation of the solvent, the residue was dissolved in ethyl acetate, and the organic phase was washed with a 5% aqueous solution of KHSO₄, with a 5% aqueous solution of NaHCO₃, and finally with brine, then was dried and evaporated, and the yellow solid obtained (600 mg) was crystallized using 1:1 isopropanol/water to obtain 450 mg of a white solid.

MS (ES⁺) : [MH⁺] = 681; HPLC (Method A2): rt = 14.7 min

i) Synthesis of cyclo{-Suc[1(R)NH₂]-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]}- (Compound A)

TFA (13 ml) was added, under stirring at 0°C, to a suspension of the compound of

Example 1(h) (400 mg) in CH₂Cl₂ (40 ml). The reaction was conducted for 2 hours at 0°C and for 2 hours at room temperature. The solvent was evaporated, the residue was treated with NaHCO₃ and water, and extracted in ethyl acetate. The organic phase was washed with brine, dried and evaporated to obtain 320 mg of a solid compound.

MS (ES⁺) : [MH⁺] = 581; HPLC (Method A2): rt = 12.4 min

A 20-mg sample was purified in an preparative HPLC to obtain 15 mg of trifluoroacetate: cyclo{-Suc[1(S)NH₂]-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂-NH]-}.TFA

MS (ES⁺): (m/z) [MH⁺] = 581; HPLC (Method A2): rt = 12.4 min; ¹H-NMR 500 MHz (DMSO): d 2.21 (dd, J = 6.1, 14.3 Hz, 1H), 2.68-2.82 (m, 5H), 2.95 (dd, J = 3.0, 14.4 Hz, 1H), 3.08 (bd, J = 12.0 Hz, 1H), 3.38 (dd, J = 3.8, 14.2 Hz, 1H), 3.48-5.36 (m, 2H), 3.98-4.08 (m, 1H), 4.11-4.17 (m, 1H), 4.20-4.28 (m, 1H), 6.71 (d, J = 9.1 Hz, 1H), 6.98 (t, J = 9.1 Hz, 1H), 7.04-7.09 (m, 2H), 7.15-7.21 (m, 4H), 7.21-7.30 (m, 6H), 7.33 (d, J = 8.1 Hz, 1H), 7.42 (d, J = 7.8 Hz), 7.67 (bs, 1H), 7.82 (bs, 1H), 8.63 (d, J = 5.2 Hz, 1H), 10.81 (d, J = 1.3 Hz, 1H)

j) Synthesis of 1-(4-piperidiny)-piperidine acetic acid hydrochloride

A solution of benzylbromo acetate (6.2 ml) in DMSO (30 ml) was added, drop by drop, to a solution of 4-piperidiny piperidine (5.0 g, titre 90%) in DMSO (50 ml) and DIPEA (4.6 ml). The mixture was stirred at room temperature for 3 hours and was then diluted with AcOEt (250 ml) and washed with water (200 ml), with a 5% aqueous solution of NaHCO₃ (3 x 100 ml), and with brine (2 x 100 ml). The pooled aqueous phases were again extracted using AcOEt (2 x 50 ml), and the pooled organic phases were dried on anhydrous sodium sulphate and evaporated to yield an orange-coloured oil, which was treated with ethyl ether (200 ml) and precipitated by addition, at 0°C, of HCl 4N in dioxane (11 ml) to obtain 7.0 g of an orange solid.

The solid was solubilized in methanol (230 ml) in the presence of a 10% Pd/C catalyst (0.5 g) and was hydrogenated at room pressure for 5 hours until complete disappearance of the reagent (control via TLC: EtOAc/i-PrOH/TEA, 80/17/3).

The reaction mixture was filtered, and the solution was dried to obtain an oil that was solidified by treatment with ethyl acetate, washed with ethyl ether, and

vacuum dried to obtain 4.7 g of a yellowish solid.

MS (ES⁺) : [MH⁺] = 227

k) HOBt (77 mg) and EDC.HCl (36 mg) were added in order to a suspension of 1-(4-piperidinyl-piperidine acetic acid hydrochloride (50 mg) in DMF (3 ml). The mixture was stirred for ten minutes, and 110 mg of Compound A were added. The clear solution was stirred for 3 hours at room temperature. The solution was dry-evaporated, and the residue was treated with acetonitrile (1 ml) and dry evaporated again. The resulting crude product was treated with AcOEt until an ivory-coloured solid (80 mg) was obtained, which was purified on preparative HPLC using the method P1 to obtain 25 mg of the final product.

MS (ES⁺) : [MH⁺] = 789.5; HPLC (Method A2): rt = 11.2 min

With a similar experimental procedure, the following compounds were obtained:

EXAMPLE 2: cyclo{-Suc[1-(R)-2(4-morpholin-4-yl-piperidin-1-yl)-acetyl amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of general formula (I) in which R = H; R₁ = H; Y = CH₂CONH; X = O)

a) Synthesis of 1-benzyloxycarbonyl-piperidin-4-one

Triethyl amine (9.2 ml) and *N*-benzyloxycarbonyl succinimide (9.0 g) were added to a solution of hydrated piperidin-4-one hydrochloride (5.0 g) in THF/H₂O (10/6, 160 ml) cooled to 5°C. The mixture was stirred for 10 minutes at 5°C and for 2 hours at room temperature.

The organic solvent was evaporated, and the resulting aqueous phase was extracted with AcOEt (3 x 70 ml). The pooled organic phases were washed with water (70 ml), with HCl 1N (70 ml), with a saturated aqueous solution of NaHCO₃ (3 x 70 ml), and with brine (70 ml). The organic phase was dried on anhydrous sodium sulphate and, after evaporation of the solvent at reduced pressure, 8.2 g of product as a viscous oil were obtained.

MS (EI⁺) : [M⁺] = 233; TLC : R_f = 0.36 (EtOAc/Hexane, 50/50); HPLC (Method A1): rt = 9.5 min

b) Synthesis of (4-piperidin-4-yl-morpholin-1-yl)-benzyloxycarbonyl

Sodium cyanoborohydride (2.39 g) and glacial acetic acid (2.80 ml) were added to a solution of 1-benzyloxycarbonyl-piperidin-4-one (8.07 g) and morpholine (2.80 ml) in methanol (140 ml), and the mixture was left to react overnight at room

temperature. The solvent was evaporated, and the residue was treated with water (70 ml) and acidified with concentrated HCl up to pH 1. The mixture was left under a suction hood until complete destruction of the reagent in excess. The resulting solution was basified by addition of solid K_2CO_3 in small portions and extracted with AcOEt (3 x 70 ml). The pooled organic phases were dried on anhydrous sodium sulphate and evaporated to yield 5.44 g of product in the form of an oil.

MS (ES^+): $[MH^+] = 305$; TLC: $R_f = 0.57$ (EtOAc/MeOH/TEA, 85/14/1); HPLC (Method A2): $rt = 8.0$ min

c) Synthesis 4-piperidin-4-yl-morpholin *bis* hydrochloride

Concentrated HCl (2.50 ml) was added to a solution of (4-piperidin-4-yl-morpholin-1-yl)-benzyloxycarbonyl (5.44 g) in methanol (150 ml), and the resulting mixture was hydrogenated at room pressure in the presence of 10% Pd/C (0.33 g) for 10 hours. The reaction mixture was filtered and washed repeatedly with methanol. The eluate was pooled and evaporated at reduced pressure. The resulting solid was washed with ethyl ether, filtered and vacuum dried to obtain 3.34 g of product.

MS (ES^+): $[MH^+] = 171$

d) Synthesis of 2-(4-morpholin-4-yl-piperidin-1-yl) acetic acid

The product was obtained according to the procedure described in point j) of Example 1, using 4-piperidin-4-yl-morpholine instead of piperidinyl piperidine.

To a solution of 2-(4-morpholin-4-yl-piperidin-1-yl) acetic acid (19 mg) in DMF (5 ml) were added HOBt (9.4 mg) and EDC.HCl (35.2 mg). After stirring for 15 minutes at room temperature, a solution of Compound A (50 mg) in DMF (5 ml) was added. The pH of the solution was corrected from pH 4 to pH 6 by addition of triethyl amine. The mixture was stirred overnight and the solvent was vacuum evaporated. The resulting residue was purified on a preparative HPLC (Method P2) to obtain 35 mg of product.

MS (ES^+): $[MH^+] = 791.5$; HPLC (Method A2): $rt = 10.9$ min

EXAMPLE 3: cyclo{-Suc[1-(R)-2(1,4')-bipiperidinyl-1'-yl-acetyl amino]-Trp-Phe-(4-F)-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of general formula (I) in which $R = F$; $R_1 = H$; $X = CH_2$; $Y = CH_2CONH$)

The compound was obtained using the procedure described in Example 1(a)-(k), but using Boc-Phe(4-F)-OH instead of Boc-Phe-OH.

MS (ES⁺): [MH⁺] = 807.4; HPLC (Method A2): rt = 11.4 min

EXAMPLE 4: cyclo{-Suc[1-(R)-2(4-morpholinyl-4-piperidinyl-1-yl-acetyl) amino]-Trp-(5-F)-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

(compound of general formula (I) in which R = H; R₁ = F; X = O; Y = CH₂CONH)

- 5 The compound was obtained using the procedure described in Example 2(a)-2(e) but using Boc-Trp(5-F)-OH instead of Boc-Trp-OH.

MS (ES⁺): [MH⁺] = 809.7; HPLC (Method A2): rt = 11.5 min

EXAMPLE 5: cyclo{-Suc[1-(R)-2(4-(1-oxo-1-thiomorpholin-4-yl-piperidin-1-yl)-acetyl amino)-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

- 10 (compound of general formula (I) in which R = H; R₁ = H; Y = CH₂CONH; X = SO)

a) Synthesis of 1-*tert*-butoxycarbonyl-4-(1-thiomorpholin-4-yl)piperidine

- To a solution of 1-*tert*-butoxycarbonyl-piperidin-4-one (2.0 g) in methanol (45 ml) were added thiomorpholin (1 ml), acetic acid (1.2 ml), and finally NaCNBH₃ (630 mg). The mixture was left to react overnight. The mixture was concentrated to a small volume, diluted with water (30 ml) and HCl 1N to pH 1.5, and extracted with ethyl acetate (3 x 20 ml). The pooled organic phases were washed with a 5% aqueous solution of NaHCO₃ and brine, and dried on anhydrous sodium sulphate. The solid was evaporated, and the resulting oil was purified on a chromatographic column, eluting with 95:5 ethyl acetate/methanol to obtain 1.4 g of product in the form of a colourless oil.

MS (ES⁺): [MH⁺] = 287; TLC: rf = 0.58 (EtOAc/MeOH, 95/5, iodine detector)

b) Synthesis of 1-oxo-1-thiomorpholin-4-yl-piperidine

- To a solution of (4-(1-thiomorpholin-4-yl-piperidin-1-yl)-*tert*-butoxycarbonyl (1.0 g) in methylene chloride (40 ml) and trifluoroacetic acid (0.3 ml) cooled to -15°C was added, drop by drop, a solution of *meta*-chloroperbenzoic acid (70%, 0.9 g) in methylene chloride (40 ml). When the addition was completed, the mixture was left to react for 30 minutes at 0°C and was diluted with dichloromethane until a homogeneous mixture was obtained. The reaction mixture was transferred into a separating funnel and washed repeatedly with a 5% aqueous solution of NaHCO₃ and brine, and was then dried on anhydrous magnesium sulphate. An oil was obtained by evaporation of the solvent.

c) The product previously obtained was solubilized in a 2:1 mixture of methylene

chloride and trifluoroacetic acid (60 ml) cooled to 5°C. The solution was left to react for 1 hour at 5°C and for 2 hours at room temperature. The reaction mixture was doubled in volume by addition of toluene, and was then vacuum evaporated until a dense oil was obtained consisting of the product in the form of trifluoroacetic salt. The amount of product obtained was 0.9 grams.

MS (ES⁺): [MH⁺] = 203

d) Synthesis of 2 (4-(1-oxo-1-thiomorpholin-4-yl)-piperidin-1-yl)-acetyl-*tert*-butyl ester

To a solution of 1-oxo-1-thiomorpholin-4-piperidine (0.9 g) in DMSO (8 ml) were added DIPEA (0.7 ml), and then *tert*-butyl-bromo acetate (0.6 ml), and the mixture was left to react overnight at room temperature. The formation of the product of reaction was revealed by TLC (chloroform/methanol 9/1, iodine detector, R_f = 0.4). The reaction mixture was diluted with ethyl acetate and washed with a 5% aqueous solution of NaHCO₃ and brine, and dried on anhydrous sodium sulphate. The solvent was evaporated, and 0.75 g of a solid was obtained, which was sufficiently pure to be used as it was.

MS (ES⁺): [MH⁺] = 317

e) Synthesis of 2(4-(1-oxo-1-thiomorpholin-4-yl)-piperidin-1-yl)-acetic acid

To a solution of 2(4-(1-oxo-1-thiomorpholin-4-yl)-piperidin-1-yl)-acetyl *tert*-butyl ester (200 mg) in dichloromethane (4 ml) was added trifluoroacetic acid (2 ml), and the solution was stirred for 3 hours. The disappearance of the reagent was verified by TLC (AcOEt/MeOH, 9:1), and the solution was diluted with toluene and dry evaporated. The resulting residue was treated with ethyl ether, and the solid that formed was filtered, washed with ether, and dried in a dryer to obtain 100 mg of product in the form of trifluoroacetate salt.

MS (ES⁺): [MH⁺] = 261

f) To a solution of 2(4-(1-oxo-1-thiomorpholin-4-yl)-piperidin-1-yl) acetic acid of the previous example (50 mg) in DMF (4 ml) were added HOBt (30 mg) and EDC.HCl (30 mg). After stirring for 15 minutes at room temperature, Compound A (50 mg) was added. The pH of the solution was corrected from pH 4 to pH 6 by addition of triethyl amine. The mixture was stirred overnight, and the solvent was vacuum evaporated. The resulting residue was purified on preparative HPLC (Method P2)

to obtain 24 mg of product.

MS (ES⁺): [MH⁺] = 824; HPLC (Method A2): rt = 10.8 min

EXAMPLE 6: cyclo{-Suc[1-(R)-2(4-(1-thiomorpholin-4-yl)-piperidin-1-yl)-acetyl amino]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

5 (Compound of general formula (I) in which R = H; R₁ = H; Y = CH₂CONH; X = S)

The compound was obtained using the procedure described in Example 5, excluding the intermediate step 5(b).

MS (ES⁺): [MH⁺] = 808; HPLC (Method A2): rt = 12.9 min

**EXAMPLE 7: cyclo{-Suc[1-(R or S)-(1,4')-bipiperidiny-1'-carbonyl]-Trp-Phe-
10 [(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (slow-moving)**

(Compound of general formula (I) in which R = H; R₁ = H; X = CH₂; Y = CO)

a) Synthesis of Boc-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH₂]

To a solution of Boc-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH-Z] (1.20 g) in methanol (36 ml) and DMF (14 ml) was added 10% Pd/C (120 mg). The mixture
15 underwent stirring and was hydrogenated at room temperature and atmospheric pressure for 2 hours. The mixture was filtered, and the solid was washed with methanol. The eluates were pooled and evaporated until a viscous oil was obtained, which was solubilized in ethyl acetate. The resulting solution was washed with water and brine, and was dried on anhydrous sodium sulphate. By
20 evaporation of the organic phase, 870 mg of a white solid were obtained.

MS (ES⁺): [MH⁺] = 584; HPLC (Method A3): rt = 11.8 min

b) Synthesis of Boc-Trp-Phe{(R)-NH-CH(CH₂-C₆H₅)-CH₂NH-[2-(4-nitro-benzyloxycarbonyl)-4-*tert*-butyl)-succin-1-yl]}

To a solution of 2-(4-nitro-benzyloxycarbonyl)-succinic acid 4-*tert*-butyl ester (424
25 mg) in DMF (20 ml) at 0°C were added HOBt (490 mg), EDCI.HCl (250 mg), and finally Boc-Trp-Phe[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH₂] (700 mg). The mixture was made to react for 2 hours at room temperature. The solvent was removed by vacuum evaporation, and the resulting residue was treated with a 5% aqueous solution of KHSO₄ to obtain a solid, which was filtered, washed with a 5% aqueous
30 solution of NaHCO₃, and with water, and was finally vacuum dried on CaCl₂ to obtain 1.05 g of a solid.

MS (ES⁺): [MH⁺] = 919; HPLC (Method A4): rt = 20.3 min

c) Synthesis of cyclo{Suc[1-(4-nitro-benzyloxycarbonyl)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-}

To 20 ml of TFA cooled to 0°C was added, in small portions, 1.0 g of Boc-Trp-Phe{(R)-NH-CH(CH₂-C₆H₅)-CH₂NH-[2-(4-nitro-benzyloxycarbonyl)-4-*tert*-butyl-succin-1-yl]}. The mixture, left to react for 30 minutes at 0°C, was vacuum concentrated and diluted with DMF, and then was evaporated again until an oil was obtained that was treated with ethyl ether to yield a solid. The solid was filtered and washed with ethyl ether until an amorphous yellow solid was obtained, consisting of H-Trp-Phe{(R)-NH-CH(CH₂-C₆H₅)-CH₂NH-[2-(4-nitro-benzyloxycarbonyl)]}-1-succinic acid; 710 mg of product were obtained.

PyBOP (160 mg) and TEA (108 µl) were added to a 200-mg solution of H-Trp-Phe{(R)-NH-CH(CH₂-C₆H₅)-CH₂NH-[2-(4-nitro-benzyloxycarbonyl)]}-1-succinic acid in DMF (10 ml) under nitrogen at 0°C. The solution underwent stirring at room temperature for hours and was monitored using HPLC. The reaction mixture was dried, and the residue was dissolved in ethyl acetate. The organic phase was washed with a 5% aqueous solution of KHSO₄, with a 5% aqueous solution of NaHCO₃, and with brine, and was finally dehydrated on anhydrous sodium sulphate. After filtration and evaporation of the solvent, a residue was obtained weighing 180 mg and consisting of the product as a mixture of diastereoisomers in position 1-succinyl. The two diastereoisomers are defined as "fast-moving" (fm) and "slow-moving" (sm).

MS (ES⁺): [MH⁺](fm) = [MH⁺](sm) = 745; HPLC (Method A3): rt(fm) = 15.1 min, rt(sm) = 15.6 min

d) To the mixture of cyclo{Suc[1-(4-nitro-benzyloxycarbonyl)]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} diastereoisomers (100 mg) was added a 1:1 mixture of water and isopropanol (3 ml) containing K₂CO₃ (34 mg). The reaction mixture was left to react for 18 hours at room temperature, and was then concentrated, diluted with water, and extracted with ethyl acetate to eliminate the non-reacted product. The aqueous phase was acidified with HCl 1N until a white opalescence was formed, and was extracted again with ethyl acetate. The organic phase of the second extraction was dried on anhydrous sodium sulphate and evaporated to yield 55 mg of a white solid.

The solid was dissolved in DMF (3 ml), and HOBt (50 mg), EDCI.HCl (32 mg), and [1,4]bipiperidine (15 μ l) were added in order. After 24 hours under stirring, the reaction mixture was diluted with 3 ml of a mixture consisting of 80:20 water/acetonitrile containing 0.1% TFA, and was purified using preparative HPLC (Method P3) to separate the two diastereoisomer products present. In this way, 30 mg of fast-moving product were obtained as a white solid.

MS (ES^+): [MH^+] = 760.4; HPLC (Method A2): rt = 12.6 min

Following a similar experimental procedure, the following compound was obtained:

EXAMPLE 8: cyclo{(-Suc[1-(S or R)-(1,4')-bipiperidiny]-1'-carbonyl]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-)} (fast-moving)

(Compound of general formula (I) in which $R = H$; $R_1 = H$; $X = CH_2$; $Y = CO$)

The compound was separated from the corresponding diastereoisomer by preparative HPLC (Method P3).

MS (ES^+): [MH^+] = 760.4; HPLC (Method A2): rt = 13.0 min

Preparative HPLC methods

Mobile phase: A = H₂O + 0.1% TFA; B = CH₃CN + 0.1% TFA

Method P1

Column: Symmetry RP18, 7 μ m, 100 Å, 19 x 300 mm

Gradient from A:B = 80:20 to A:B = 50:50 in 60 min, then from A:B = 50:50 to A:B = 20:80 in 120 min

Flow rate: 15 ml/min

λ = 220, 270 nm

Method P2

Column: Symmetry RP18, 7 μ m, 100 Å, 19 x 300 mm

Gradient from A:B = 80:20 to A:B = 20:80 in 120 min

Flow rate: 15 ml/min

λ = 220, 270 nm

Method P3

Column: Vydac RP18, 20 μ m, 22 x 250 mm

Gradient from A:B = 80:20 to A:B = 20:80 in 120 min

Flow rate: 20 ml/min

λ = 240 nm

Analytical HPLC methods

Mobile phase: A = H₂O + 0.1% TFA; B = CH₃CN + 0.1% TFA

Method A1

Column: Symmetry C18, 5 mm, 100 Å, 3.9 x 150 mm

5 Gradient from A:B = 80:20 to A:B = 20:80 in 20 min

Flow rate: 1 ml/min

I = 215 nm

Method A2

Column: Luna 5 µm, C8(2), 100 Å, 4.6 x 250 mm

10 Gradient from A:B = 80:20 to A:B = 20:80 in 20 min

Flow rate: 1 ml/min

I = 220, 270 nm

Method A3

Column: Symmetry C8 5 mm, 100 Å, 3.9 x 150 mm

15 Gradient from A:B = 80:20 to A:B = 20:80 in 20 min

Flow rate: 1 ml/min

I = 220, 270 nm

Method A4

Column: Symmetry C8 5 mm, 100 Å, 3.9 x 150 mm

20 Gradient from A:B = 80:20 to A:B = 20:80 in 20 min followed by A:B = 20:80 for 6 min

Flow rate: 1 ml/min

I = 220, 270 nm

Abbreviations: For the nomenclature and abbreviations of the amino acids,
25 reference is made to the recommendations of IUPAC-IUB Joint Commission on Biochemical Nomenclature (Eur. J. Biochem. 1984, 138, 9); the amino acids are understood as being in the S configuration if not otherwise specified. The other abbreviations used are: Bzl = benzyl; DIEA = DIPEA = *N,N*-diisopropyl ethyl amine; DMF = dimethyl formamide; EDC = EDCI = 1-(3-dimethylaminopropyl)3-ethylcarbodiimide;
30 PyBOP = benzotriazol-1-yl-oxy-*tris*-pyrrolidino-phosphonohexafluorophosphate; TEA = triethyl amine; TFA = trifluoroacetic acid; Z = Cbz = *N*-benzyloxycarbonyl; Boc = *tert*-butoxycarbonyl; -Suc- = succinyl; DMF

= *N,N*-dimethyl formamide; NKA = neurokinin A; HOBt = 1-hydroxybenzotriazole; rt = retention time; THF = tetrahydrofuran. The numbering of the substituents on the succinic group designated as -Suc(1-NH₂)- is obtained with Y1 = NH₂.

Biological Activity

5 The compounds described in the present invention act as antagonists to the NK2 receptor of tachykinins. The biological activity was evaluated in three *in vitro* functional tests, using rabbit pulmonary artery (RPA), hamster trachea (HT) and rat urinary bladder (RUB), according to the methods described by C.A. Maggi *et al.*, Br. J. Pharmacol., 1990, 100, 588; P. D'Orleans-Juste *et al.*, Eur. J. Pharmacol., 1986, 125, 37; and C.A. Maggi *et al.*, J. Pharmacol. Exp. Ther., 1988, 246, 308. The affinity of the compounds for the human NK2 receptor was evaluated in a binding test using membranes of Chinese hamster ovary (CHO) cells transfected with the NK2 receptor of human ileum and the radioligand [¹²⁵I]NKA (Amersham, non-specific activity 2000 Ci/mmol) at the concentration of 100 pM in competition studies. The substances under examination were tested in a concentration range between 0.01 nM and 10 mM. At the end of incubation (30 min, 20°C) the test specimens were filtered, and radioactivity was determined using a gamma-counter.

The data emerging from the functional studies were expressed as pA₂ (O. Arunlakshana and H.O. Schild, Br. J. Pharmacol. Chemother., 1959, 14, 45), and the data emerging from the binding studies were expressed as pKi (log Ki calculated with the LIGAND program; P.J. Munson *et al.*, Anal. Biochem., 1980, 107, 220).


The compounds of the invention were found to be active in the tests referred to above, with pA₂ values of up to 9.8 and pKi values of up to 10.3.

Table of activity

Compound	pKi	pA ₂		
(example)		RPA	HT	RUB
WO9834949; Ex. 27	8.5	7.8	8.5	
WO9834949; Ex. 35	8.6	8.4	8.5	
WO9834949; Ex. 36	8.7	7.9		
WO9834949; Ex. 37	8.8			8.2
WO9834949; Ex. 46	8.1	7.7	8.3	7.7
Example 1	9.4	8.7	8.8	9.2
Example 2	9.7			9.8
Example 3	9.6			9.3
Example 4	10.1			9.7
Example 7	10.3			9.1
Example 8	9.5			9.1

CLAIMS

1. Compounds having general formula (I):



(1)

in which:

X may be CH₂, O, S, or SO;

Y is $\text{CH}_2\text{-CO-NH}$ or CO ;

R and R₁, same or different from one another, represent an H or halogen group chosen from among fluorine, chlorine, bromine, and iodine, including the respective diastereoisomers and their mixtures, with the proviso that:

when Y is $\text{CH}_2\text{-CO-NH}$, X is CH_2 , and R and R_1 are H, then the stereochemistry of the carbon atom linked to Y is of the R type;

and the pharmaceutically acceptable salts of the compounds of general formula (I) with organic and inorganic acids chosen from among the group comprising hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, carbonic acid, acetic acid, trifluoroacetic acid, trichloroacetic acid oxalic acid, malonic acid, malic acid, succinic acid, tartaric acid, citric acid, methanesulphonic acid, and p-toluenesulphonic acid.

2. Compounds according to Claim 1 in which:


R and R₁ may be the same as or different from one another and are chosen from the group consisting of H or F, and the other substituents are as defined above.

3. Compounds according to Claim 2, as listed below:

- 2 i) cyclo{-Suc[1-(R)-2(1,4')-bipiperidinyl-1'-yl-acetyl amino]-Trp-Phe-[(R)-NH-
3 CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = CH₂)
- 4 ii) cyclo{-Suc[1-(R)-2(4-morpholin-4-yl-piperidin-1-yl)-acetyl amino]-Trp-Phe-[(R)-
5 NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = O)
- 6 iii) cyclo{-Suc[1-(R)-2(1,4')-bipiperidinyl-1'-yl-acetyl amino]-Trp-Phe-(4-F)-[(R)-NH-
7 CH(CH₂-C₆H₅)-CH₂NH]-} (R = F; R₁ = H; Y = CH₂CONH; X = CH₂)
- 8 iv) cyclo{-Suc[1-(R)-2(4-morpholinyl-4-piperidinyl-1-yl-acetyl) amino]-Trp-(5-F)-
9 Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = F; Y = CH₂CONH; X = O)
- 10 v) cyclo{-Suc[1-(R)-2(4-(1-oxo-1-thiomorpholin-4-piperidin-1-yl)-acetyl amino]-Trp-
11 Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = SO)
- 12 vi) cyclo{-Suc[1-(R)-2(4-(1-thiomorpholin-4-yl-piperidin-1-yl)-acetyl amino]-Trp-
13 Phe-[(R)-NH-CH(CH₂-C₆H₅)-CH₂NH]-} (R = H; R₁ = H; Y = CH₂CONH; X = S)
- 14 vii) cyclo{-Suc[1-(1,4')-bipiperidinyl-1'-carbonyl]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-
15 CH₂NH]-} (slow-moving) (R = H; R₁ = H; Y = CO; X = CH₂)
- 16 viii) cyclo{-Suc[1-(1,4')-bipiperidinyl-1'-carbonyl]-Trp-Phe-[(R)-NH-CH(CH₂-C₆H₅)-
17 CH₂NH]-} (fast-moving) (R = H; R₁ = H; Y = CO; X = CH₂).

1 4. Process for the preparation of compounds having general formula (I)
2 characterized by the following steps of synthesis:


- 3 i) combining the diamine system represented by (R)-1-benzyl-2-(N-
4 benzyloxycarbonylamino)-ethyl amine, according to methodologies known in the
5 chemistry of peptides, with Phe and Trp, possibly substituted, respectively by an R
6 group and an R₁ group, in which said groups have the meanings previously
7 defined;
- 8 ii) condensing to the above the dicarboxylic system represented by a derivative of
9 succinic acid, and cyclizing the linear product obtained to yield the desired
10 monocyclic system having general formula (III)



(III)

in which Y_1 may be $-NH_2$ or $-COOH$, and R and R_1 have the meanings described above.

iii) finally condensing the compounds of Formula (III) with the compounds of Formula (IV)



(IV)

in which Y_2 may be H or CH_2-COOH , and X has the meaning previously defined, to yield the final compounds of general formula (I).

5. Pharmaceutical compositions containing as active principle compounds having general formula (I) according to Claims 1-3 in combination with pharmaceutically acceptable carriers or excipients.

6. Pharmaceutical compositions according to Claim 5 for use as tachykinin antagonists.

7. Pharmaceutical compositions according to Claim 6 for use as antagonists towards the human NK2 receptor.

8. Pharmaceutical compositions according to Claim 7 for use in the treatment of the bronchospastic component of asthma, coughing, pulmonary irritations, intestinal spasms or local spasms of the bladder and the ureter during cystitis, renal infections and colics.

9. Use of a compound according to Claims 1-3 for the preparation of pharmaceutical compositions to be employed to antagonize the effects of

- 3 tachykinins in illnesses in which these play an important role.
- 1 10. Use of a compound according to Claim 9 in which NK2 is specifically
2 antagonized.
- 1 11. Use of a compound according to Claims 9 and 10 for the treatment of the
2 bronchospastic component of asthma, coughing, pulmonary irritations, intestinal
3 spasms or local spasms of the bladder and the ureter during cystitis, renal
4 infections and colics.
- 1 12. Method for the treatment of the bronchospastic component of asthma,
2 coughing, pulmonary irritations, intestinal spasms or local spasms of the bladder
3 and the ureter during cystitis, renal infections and colics, in which the patient is
4 administered between 0.1 and 10 mg/kg of body weight of active principle
5 consisting of products of Formula (I) according to Claim 1.

THIS PAGE BLANK (USPTO)


(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
26 April 2001 (26.04.2001)

PCT

(10) International Publication Number
WO 01/29066 A3

- (51) International Patent Classification⁷: C07K 5/02. (74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi S.p.A., Corso dei Porta Vittoria, 9, I-20122 Milano (IT).
5/06, 1/06, A61K 38/05, A61P 11/06
- (21) International Application Number: PCT/EP00/10181 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 17 October 2000 (17.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: FI99A000217 21 October 1999 (21.10.1999) IT (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): MENARINI RICERCHE S.P.A. [IT/IT]; Via Tito Speri, 10, I-00040 Pomezia (IT).
- (72) Inventors; and (75) Inventors/Applicants (*for US only*): PERROTTA, Enzo [IT/IT]; Via Ragazzi del '99, 76, I-50141 Firenze (IT). GIANNOTTI, Danilo [IT/IT]; Via Roma, 128, I-55011 Altopascio (IT). ALTAMURA, Maria [IT/IT]; Viale Matteotti, 54, I-50132 Firenze (IT). GIULIANI, Sandro [IT/IT]; Via Carlo Borroni, 11, I-50012 Bagno a Ripoli (IT). MAGGI, Carlo, Alberto [IT/IT]; Via Michelazzi, 43, I-50141 Firenze (IT).
- Published:
— with international search report
- (88) Date of publication of the international search report:
1 November 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: BASIC MONOCYCLIC COMPOUNDS HAVING NK2 ANTAGONIST ACTION, PROCESSES FOR THEIR PREPARATION, AND FORMULATIONS CONTAINING THEM



(57) Abstract: Basic monocyclic compounds having general formula (I) useful as NK-2 antagonists are described, processes for their preparation and pharmaceutical compositions containing them are also described.

WO 01/29066 A3

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

Internat'l Application No.
PCT/EP 00/1

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07K5/02 C07K5/06 C07K1/06 A61K38/05 A61P11/06		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, CHEM ABS Data, BIOSIS, EMBASE, MEDLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 00 08046 A (MAGGI CARLO ALBERTO ;MENARINI RICERCHE SPA (IT); GUIDI ANTONIO (IT) 17 February 2000 (2000-02-17) the whole document ---	1-12
X	WO 98 34949 A (BUGNO CRISTINA DI ;MAGGI CARLO ALBERTO (IT); MENARINI RICERCHE SPA) 13 August 1998 (1998-08-13) cited in the application claim 3; examples 1,46 ---	1-12
A	WO 93 03059 A (MENARINI FARMA IND) 18 February 1993 (1993-02-18) ---	
A	EP 0 333 174 A (FUJISAWA PHARMACEUTICAL CO) 20 September 1989 (1989-09-20) --- -/--	
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents :</p> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*G* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">28 March 2001</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">03/05/2001</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Cervigni, S</div>

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/10181

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 17362 A (BOEHRINGER INGELHEIM KG) 15 May 1997 (1997-05-15) -----	

INTERNATIONAL SEARCH REPORT

Internat'l Appl. No.

PCT/EP 00/10181

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0008046 A	17-02-2000	IT F1980186 A AU 5507999 A	07-02-2000 28-02-2000
WO 9834949 A	13-08-1998	IT F1970020 A AU 6620198 A CN 1246866 T EP 1017711 A ZA 9800931 A	07-08-1998 26-08-1998 08-03-2000 12-07-2000 07-08-1998
WO 9303059 A	18-02-1993	IT 1258939 B IT 1251164 B AU 2387592 A EP 0606222 A JP 6509571 T PT 100764 A	08-03-1996 04-05-1995 02-03-1993 20-07-1994 27-10-1994 28-02-1994
EP 0333174 A	20-09-1989	AT 137763 T AU 3132489 A CA 1329444 A CN 1037156 A DE 68926403 D DE 68926403 T DK 126389 A FI 891176 A HU 49628 A JP 1287095 A NO 891082 A US 5187156 A ZA 8901551 A	15-05-1996 21-09-1989 10-05-1994 15-11-1989 13-06-1996 17-10-1996 17-09-1989 17-09-1989 30-10-1989 17-11-1989 18-09-1989 16-02-1993 29-11-1989
WO 9717362 A	15-05-1997	DE 19541283 A AU 7562296 A EP 0871653 A JP 2000501382 T ZA 9609248 A	07-05-1997 29-05-1997 21-10-1998 08-02-2000 06-05-1997

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)